



REFERENCES

- Anet, F.A.L., Hughes, G.K., and Ritchie, E. 1952. Alkaloids of Australian *Strychnos* species. I. Constituents of *S. lucida* R.Br. and *S. psilosperma* F.Muell. Aust.J.Chem. 6 : 58-65.
- Angenot, L., Dubois, M., Ginion, Ch., Dorsser, W. van, and Dresse, A. 1975. Chemical structure and pharmacological (curarizing) properties of various indole alkaloids extracted from an African *Strychnos*. Arch.int.Pharmacodyn. 215 : 246-258.
- Baser, K.H.C. 1978. Alkaloids of Strychnos nux-vomica. Ph.D. dissertation. University of London.
- Bassleer, R., et al. 1982. Effect de trois alcaloïdes extraits du *Strychnos usambarensis* sur des cellules cancéreuses en culture. Planta Medica 45 : 123-126.
- Bassleer, R., et al. 1983. Etude complémentaire de la cytotoxicité de la mélinonine F, alcaloïde dérivé de la β-carboline. Planta medica 49 : 158-161.
- Battersby, A.R., and Hall, E.S. 1969. The intermediacy of geissoschizine in indole alkaloid biosynthesis : Rearrangement to the *Strychnos* skeleton. Chem.Comm. : 793-794.

- Battersby, A.R., Brown, S.H., and Payne, T.G. 1970. Bio-synthesis of loganin and the indole alkaloids from hydroxygeraniol-hydroxynerol. Chem. Comm. :827-828.
- Battersby, A.R., Burnett, A.R., and Parsons, P.G. 1969a. Alkaloid biosynthesis. Part XIV. Secologanin : Its conversion into ipecoside and its role as biological precursor of the indole alkaloids. J. Chem. Soc. (C) : 1187-1192.
- _____. 1969b. Alkaloid biosynthesis. Part XV. Partial synthesis and isolation of vincoside and isovincoside : Biosynthesis of the three major classes of indole alkaloids from vincoside. J. Chem. Soc. (C) : 1193-1200.
- _____. 1970. Preparation and isolation of deoxyloganin : Its role as precursor of loganin and the indole alkaloids. Chem. Comm. : 826-828.
- Battersby, A.R., Hall, E.S., and Southgate, R. 1969. Alkaloid biosynthesis. part XIII. The structure, stereochemistry, and biosynthesis of loganin. J. Chem. Soc. (C) : 721-728.
- Battersby, A.R., Kapil, R.S., Martin, J.A., and Mrs. Lucy Mo. 1968. Loganin as precursor of the indole alkaloids. Chem. Comm. : 133-134.

Battersby, A.R., Brown, R.T., Kapil, R.S., Martin, J.A., and Plunkett, A.O. 1966. Role of loganin in the biosynthesis of indole alkaloids. Chem.Comm. : 890-891.

Battersby, A.R., Brown, R.T., Knight, J.A., Martin, J.A., and Plunkett, A.O. 1966. Biosynthesis of the indole alkaloids from a monoterpenone. Chem.Comm. : 346-347.

Battersby, A.R., Byrne, J.C., Kapil, R.S., Martin, J.A., and Payne, T.G. 1968. The mechanism of indole alkaloid biosynthesis. Chem.Comm. : 951-953.

Baxter, C., and Slaytor, M. 1972. Partial purification and some properties of tryptophan decarboxylase from *Phalaris tuberosa*. Phytochemistry 11 : 2763-2766.

Beljanski, M., and Bugiel, J. 1979. Demande française de brevet d'invention no. 7807155. quoted in Ohiri, F.C., Verpoorte, R., and Baerheim Svendsen, A. 1983. the African *Strychnos* species and their alkaloids : A review. J. of Ethnopharmacology 9 : 167-223.

Bernauer, K., Karrer, P., and Schmid, H. 1963. C-Dihydrotoxiferine dehalides and a process for their preparation. U.S. patent no. 3,073,832. quoted in Ohiri, F.C., Verpoorte, R., and Baerheim

- Svendsen, A. 1983. the African *Strychnos* species and their alkaloids : A review. J. of Ethnopharmacology 9 : 167-223.
- Biemann, K., Grossert, J.S., Occolowitz, J., and Warren, F.L. 1965. The indole alkaloids. Part V. The structure of henninsoline. J.Chem.Soc. : 2818-2822.
- Bisset, N.G. 1966. The arrow and dart poisons of South-East Asia, with particular reference to the *Strychnos* species used in them. Part I. Indonesia, Borneo, Philippines, Hainan, and Indo-china. Lloydia 29 : 1-18.
- _____. 1970. The African species of *Strychnos*. Part I. The ethnobotany. Lloydia 33 : 201-243.
- _____. 1972a. Chemical studies on the alkaloids of Asian and African *Strychnos* species. Lloydia 35 : 203-206.
- _____. 1972b. The Asian species of *Strychnos*. Part I. *Strychnos* as a source of drug Lignum colubrinum (snake-wood). Lloydia 35 : 95-116.
- _____. 1974. The Asian species of *Strychnos*. Part III. The ethnobotany. Lloydia 37 : 62-107.
- _____. 1980. Alkaloids of the Loganiaceae. In J.D. Phillipson, and M.H. Zenk (eds.), Indole and

biologically related alkaloids. pp.27-61. London : Academic Press.

Bisset, N.G., and Khalil, A.A. 1976. New alkaloids from *Strychnos icaja*. Phytochemistry 15 : 1973-1976.

Bisset, N.G., and Leeuwenberg, A.J.M. 1968. The use of *Strychnos* species in Central African ordeal and arrow poisons. Lloydia 31 : 208-222.

Bisset, N.G., and Phillipson, J.D. 1971. the African species of *Strychnos*. Part II. The alkaloids. Lloydia 34 : 1-60.

_____. 1973. Alkaloids from the seeds of *Strychnos wallichiana* Steud. ex DC. (*Strychnos cinnamomifolia* Thwaites var. *wightii* A.W.Hill). J.Phar.Pharmac. 25 : 563-569.

_____. 1976. The Asian species of *Strychnos*. Part IV. The alkaloids. Lloydia 39 : 263-325.

Bisset, N.G., and Woods, M.C. 1966. The arrow and dart poisons of South-East Asia, with particular reference to the *Strychnos* species used in them. Part II. Burma, Thailand, and Malaya. Lloydia 29 : 172-195.

Bisset, N.G., Baser, K.H.C., Phillipson, J.D., Bohlin, L., and Sandberg, F. 1977. Muscle-relaxant activity in Asian *Strychnos* species. A re-examination of

two Western Malaysian dart poisons. Lloydia 40 : 546-560.

Bisset, N.G., et al. 1973. The Asian species of *Strychnos*. Part II. Typification, miscellaneous notes, synoptic key, and sectional classification. Lloydia 36 : 179-201.

Bohlin, L. 1978. Some Strychnos alkaloids : Their occurrence, structure and biological activity. Ph.D dissertation. Uppsala University.

Bohlin, L., and Iskander, G.M. 1978. Muscle-relaxant activity of 2-carboxamidostrychnine derivatives. Acta Pharm. Suec. 16 : 41-46.

Bohlin, L., Ali, Y., and Iskander, G.M. 1975. Synthesis and convulsant effect of 2- and 3-substituted strychnine derivatives. Acta Pharm. Suec. 12 : 461-466.

Bohlin, L., Ali, Y., and Sandberg, F. 1974. Screening of African *Strychnos* species for convulsant and muscle-relaxant effects. Acta Pharm. Suec. 11 : 233-238.

Bohlin, L., Rolfsen, W., Strömbom, J., and Verpoorte, R. 1979. Alkaloids and biological activity of *Strychnos angolensis*. Planta medica 35 : 19-30.

- Borstel, K.von, and Hartmann, T. 1986. Selective uptake of pyrrolizide *N*-oxides by cell suspension cultures from pyrrolizidine alkaloid producing plants. Plant Cell Reports 5 : 39-42.
- Brown, R.T., Leonard, J., and Sleigh, S.K. 1978. The role of strictosidine in monoterpenoid indole alkaloid biosynthesis. Phytochemistry 17 : 899-900.
- Caprasse, M., and Angenot, L. 1982. Cytotoxic anhydronium bases from *Strychnos usambarensis*. Planta Medica 45 : 149.
- Caron, C., et al. 1988. Antimicrobial and antifungal activities of quasi-dimeric and related alkaloids. Planta Medica 54 : 409-412.
- Charoendee Pingsuthiwong. 1986. Alkaloids from the stem of *Strychnos ignatii* Berg. Master's Thesis. Chulalongkorn University.
- Cordell, G.A. 1974. The biosynthesis of indole alkaloids. Lloydia 37 : 219-298.
- Coscia, C.J., Botta, L., and Guarnaccia, R. 1970. On the mechanism of iridoid and secoiridoid monoterpenone biosynthesis. Archives of Biochemistry and Biophysics 136 : 498-506.
- Coune, C. 1980. Contribution à l'étude du *Strychnos gossweileri* du Zaïre : Isolement et détermination

de structure de nouveaux alcaloïdes. Ph.D dissertation. Universite de Liege. quoted in Ohiri, F.C., Verpoorte, R., and Baerheim Svendsen, A. 1983. The African *Strychnos* species and their alkaloids : A review. J. of Ethnopharmacology 9 : 167-223.

Craib, W.G., and Kerr, A.F.G. 1951. Florae Siamensis enumeratio. A list of the plants known from Siam with records of their occurrence (vol.III, part I). Bangkok : The auspecies of the Siam society.

Dubois, M., Ginion, C., Angenot, L., Van Dresser, W., and Dresse, A. 1974. Charactères chimiques et propriétés pharmacologiques de plusieurs alcaloïdes extraits de *Strychnos usambarensis*. Archives Internationales de Physiologie de Biochimie 82 : 823-830. quoted in Ohiri, F.C., Verpoorte, R., and Baerheim Svendsen, A. 1983. The African *Strychnos* species and their alkaloids : A review. J. of Ethnopharmacology 9 : 167-223.

Escher, S., Loew, P., and Arigoni, D. 1970. The role of hydroxygeraniol and hydroxynerol in the biosynthesis of loganin and indole alkaloids. Chem. Comm. : 823-825.

Goonetilleke, A., Rolfsen, W., and Rajapakse, L. 1980. Tertiary indole alkaloids of *Strychnos aculeata*. Planta Medica 39 : 208.

- Grossert, J.S., Hugo, J.M., von Klemperer, M.E., and Warren, F.L. 1965. The indole alkaloids. Part III. The isolation of diaboline and three new alkaloids, henningsamine, henningsoline and rindline from *Strychnos henningsii* Gilg. J.Chem.Soc. : 2812-2814.
- Guarnaccia, R., and Coscia, C.J. 1971. Occurrence and biosynthesis of secologanic acid in *Vinca rosea*. J.A.C.S. 93 : 6320-6321.
- Hall, E.S., McCapra, F., and Money, T. 1966. Concerning the terpenoid origin of indole alkaloids : Biosynthetic mapping by direct mass spectrometry. Chem.Comm. : 348-350.
- Heimberger, S.I., and Scott, A.I. 1973. Biosynthesis of strychnine. J.C.S.Chem.Comm. : 217-219.
- Herbert, R.B. 1981a. Biosynthesis. In M.F. Grundon (ed.), The alkaloids, specialist periodical reports. vol.10, pp.19-22. London : The Royal Society of Chemistry.
- _____. 1981b. Biosynthesis. In M.F. Grundon (ed.), The alkaloids, specialist periodical reports, vol.11, pp.17-20. London : The Royal Society of Chemistry.

Hesse, M. 1974. Indolalkaloide, Teil 1 : Text, Progress in mass spectrometry. vol.1, pp.85-90. Germany : Verlag Chemie.

Hill, A.W. 1930. *Strychnos silvicola* A.W.Hill. Kew. Bull. : 156.

Hokanson, G.C. 1976. Potential antitumor agents from higher plants. Part I : *Strychnos henninsii*. Part II : *Centaurea solstitialis*. Part III : *Croton texensis*. Dissertation Abstract International Section B 37, no.3 : 1265-B.

Inouye, H., Ueda, S., Aoki, Y., and Takeda, Y. 1972. Studies on monoterpenoid glucosides and related natural products. XVII. The intermediacy of 7-desoxyloganic acid and loganin in the biosynthesis of several iridoid glucosides. Chem.Pharm.Bull. 20 : 1287-1296.

Iskander, G.M., and Bohlin, L. 1978. Structure-activity relationship of strychnine derivatives modified in the non-aromatic part. Acta Pharm.Suec. 15 : 431-438.

Kan-Fan, C., and Husson, H.-P. 1979. Isolation and biomimetic conversion of 4,21-dehydrogeissoschizine. J.C.S.Chem.Comm. : 1015-1016.

Karrer, von P., Eugster, C.H., and Waser, P. 1949. Über curarewirkung einiger strychnidin-und dihydro-



strychnidin-chlor-alkylate. Helv.Chim.Acta 32 : 2381-2385.

Kisakürek, M.V., and Hesse, M. 1980. Chemotaxonomy studies of the Apocynaceae, Loganiaceae and Rubiaceae, with reference to indole alkaloids. In J.D. Phillipsin, and M.H. Zenk (eds.), Indole and biologically related alkaloids. pp.11-26. London : Academic Press.

Kisakürek, M.V., Leeuwenberg, A.J.M., and Hesse, M. 1983. A chemotaxonomy investigation of the plant families of Apocynaceae, Loganiaceae, and Rubiaceae by their indole alkaloid content. In S.W. Pelletier (ed.). Alkaloids : Chemical and biochemical perspectives, volume 1, pp.211-376. New York-Chichester-Brishane-Toronto-Singapore : John Wiley & Son.

Kompiš, I., Hesse, M., and Schmid, H. 1971. An approach to the biogenetic classification of indole alkaloids. Lloydia 34 : 269-291.

Krukoff, B.A. 1972. American species of *Strychnos*. Lloydia 35 : 193-194.

Leclercq, J., Pauw-Gillet, M.-C. de, Bassleer, R., and Angenot, L. 1986. Screening of cytotoxic activities of *Strychnos* alkaloids (method and result). J. of Ethnopharmacology 15 : 305-316.

Leenhouts, P.W. 1962. Loganiaceae. In C.G.G.J. van Steenis (ed.), Flora Malesiana (ser.I, vol.6), pp.354-356. Groningen, Netherlands : Wolters-Noordhoff.

Leeuwengberg, A.J.M. 1969. The Loganiaceae of Africa. VIII. *Strychnos* III. (Revision of the African species with notes on the extra-African). Mededel. Landbphgsch. Wageningen 69 : 1-316. quote in Bisset, N.G., and Phillipson, J.D. 1971. The African species of *Strychnos*. Part II. The alkaloids. Lloydia 34 : 1-60.

_____. 1980. The taxonomy position of some genera in the Loganiaceae, Apocynaceae and Rubiaceae, related families which contain indole alkaloids. In J.D. Phillipson, and M.H. Zenk (eds.), Indole and biologically related alkaloids, pp.1-10. London : Academic Press.

Leonard, B.E. 1965a. An investigation of the pharmacology of macusine B. J.Pharm.Pharmac. 17 : 566-576.

_____. 1965b. Macusine B : further pharmacology. J.Pharm.Pharmac. 17 : 788-793.

Lin, R.C.Y. 1952. Pharmacological and toxicological actions of dart poison used in Malaya. Nature 170 : 416.

- Loew, P., and Arigoni, D. 1968. The biological conversion of loganin into indole alkaloids. Chem. Comm. : 137-138.
- Loew, P., Goeggel, H., and Arigoni, D. 1966. A monoterpenoid precursor in the biosynthesis of indole alkaloids. Chem. Comm. : 347-348.
- Marini-Bettolo, G.B. 1959. Curarizing alkaloids of *Strychnos*. In D. Bovet, F. Bovet-Nitti, and G.B. Marini-Bettolo (eds.), Curare and curare-like agents. pp.137-148. Amsterdam-London-New York-Princeton : Elsevier Publishing Company.
- Marini-Bettolo, G.B., and Bisset, N.G. 1972. Chemical studies on the alkaloid of American *Strychnos* species. Lloydia 35 : 195-202.
- Marini-Bettolo, G.B., Ciaca, M.A., Galeffi, C., Bisset, N.G., and Krukoff, B.A. 1972. The occurrence of strychnine and brucine in an American species of *Strychnos*. Phytochemistry 11 : 381-384.
- Massiot, G., et al. 1983. Occurrence of longicaudatine, a new type of bis-indole base and bisnor-C-alkaloid H in *Strychnos* species. J.Org.Chem. 48 : 1869-1872.
- _____. 1988. Alkaloids from roots of *Strychnos matopensis*. Phytochemistry 27 : 3293-3304.

- Melo, M. de F.F., Santos, C.A. de M., Chiappeta, A, de A., Mello, J.F. de, and Mukherjee, R. 1987. Chemistry and pharmacology of a tertiary alkaloid from *Strychnos trinervis* root bark. J. of Ethnopharmacology 19 : 319-325.
- Nagakura, N., Rueffer, M., and Zenk, M.H. 1979. The biosynthesis of monoterpenoid indole alkaloids from strictosidine. J.C.S.Perkin I : 2308-2312.
- Neuss, N. 1980. The spectrum of biological activities of indole alkaloids. In J.D. Phillipson, and M.H. Zenk (eds.), Indole and biologically related alkaloids, pp.293-313. London : Academic Press.
- Ohiri, F.C., Verpoorte, R., and Baerheim Svendsen, A. 1983. The African *Strychnos* species and their alkaloids : A review. J. of Ethnopharmacology 9 : 167-223.
- _____. 1984. 12-Hydroxy-11-methoxy-diaboline : A new alkaloid from *Strychnos spinosa*. Planta Medica 50 : 446-447.
- Olaniyi, A., Rolfsen, W., and Verpoorte, R. 1981. Quaternary indole alkaloids of *Strychnos decussata*. Planta medica 43 : 353-359.
- Popjak, G. 1970. Conversion of mevalonic acid into phenyl hydrocarbons as exemplified by the

- synthesis of squalene. In T.W. Goodwin (ed.), Natural substances formed biologically from mevalonic acid, pp.17-33. London : Academic Press.
- Rapepol Bavovada. 1983. Alkaloids of Asian *Strychnos* species. Ph.D. dissertation. University of London.
- Renrong, L., and Lida, L. 1985. Alkaloids of *Strychnos cathayensis* Merr. Huaxue Xuebao 43 : 775-7. Chemical Abstracts 103 : 157408g.
- Rolfsen, W., Olaniyi, A., and Hylands, P. 1980. New tertiary alkaloids of *Strychnos decussata*. Journal of Natural Products 43 : 97-102.
- Rolfsen, W., Olanyi, A., and Sandberg, F. 1980. Muscle-relaxant activity of decussine, a new indole alkaloid of *Strychnos decussata*. Acta Pharm. Suec. 17 : 105-111.
- Rolfsen, W., Hakizadeh, Z.M., Sandberg, F., and Strömbom, J. 1978. Pharmacological and phytochemical screening of *Strychnos* species from Gabon. Acta Pharm. Suec. 16 : 47-55
- Rolfsen, W., Bohlin, L., Yeboah, S.K., Geevaratne, M., and Verpoorte, R. 1978. New indole alkaloids of *Strychnos dale* and *Strychnos elaeocarpa*. Planta Medica 34 : 264-273.

Rolfsen, W., Olaniyi, A., Verpoorte, R., and Bohlin, L. 1981. Some new decussine-type alkaloids from *Strychnos decussata*, *Strychnos dale* and *Strychnos elaeocarpa*. Journal of Natural Products 44 : 415-421.

Rolfsen, W., et al. 1980. Isolation, synthesis and pharmacological activity of strychnocarpine, a new β -carboline alkaloid from *Strychnos elaeocarpa*. Acta Pharm.Suec. 17 : 333-340.

Rüeffer, M., Nagakura, N., and Zenk, M.H. 1978. Stric-tosidine, the common precursor for monoterpenoid indole alkaloids with 3α and 3β configuration. Tet.Let. 18 : 1593-1596.

Rüeffer, M., Kan-Fan, C., Husson H.-P., Stöckigt, J., and Zenk, M.H. 1979. 4,21-Dehydrogeissoschizine, an intermediate in heteroyohimbine alkaloid biosynthesis. J.C.S.Chem.Comm. : 1016-1018.

Schlatter, von Ch., Waldner, E.E., Schmid, H., Maier, W., and Gröger, D. 1969. Zur biosynthese des strychnins. Helv.Chim.Acta 52 : 776-789.

Scott, A.I., and Lee, S.-L. 1975. Biosynthesis of the indole alkaloids. A cell-free system from *Catharanthus roseus*. J.A.C.S. : 6906-6908.

Scott, A.I., Cherry, P.C., and Qureshi, A.A. 1969. Mechanism of indole alkaloid biosynthesis. The

corynanthe-strychnos relationship. J.A.C.S. 91 :
4932-4933.

Singh, H., and Kapoor, K.V. 1976. Investigation of *Strychnos* spp. IV. Pharmacological studies of alkaloids of *Strychnos potatorum*. Planta Medica 29 : 226-233.

Southon, I.W., and Buckingham, J. 1989. Dictionary of alkaloids. London, New York : Chapman and Hall.

Stöckigt, J. 1980. The biosynthesis of heteroyohimbine-type alkaloids. In J.D. Phillipson, and M.H. Zenk (eds.), Indole and biologically related alkaloids. pp.113-1416. London : Academic Press.

Stöckigt, J., and Zenk, M.H. 1977. Strictosidine (isovincoside) : the key intermediate in the biosynthesis of monoterpenoid indole alkaloids. J.C.S.Chem.Comm. : 646-648.

Stöckigt, J., Hofle, G., and Pfitzner, A. 1980. Mechanism of the biosynthetic conversion of geissoschizine to 19-epi-ajmalicine in *Catharanthus roseus*. Tet.Let. 21 : 1925-1926.

Tem Smitinand, 1980. Thai plant names (Botanical names-vernacular names). 2nd ed., pp.319-320. Bangkok : Funny publishing.

Thepenier, P., Jacquier, M.-J., Massiot, G., Le Men-Oliver, L., and Delaude, C. 1988. Alkaloids from *Strychnos staudtii*. Phytochemistry 27 : 657-659.

Thomas, R. 1961. A possible biosynthetic relationship between the cyclopentanoid monoterpenes and the indole alkaloids. Tet.Let. 16 : 544-553.

Tit, M. 1982. Contribution à l'étude du Strychnos variabilis, plante toxique du Bas-Zaire : isollement et détermination de structure de nouveaux alcaloïdes. Ph.D. dissertation. Université de Liège. quote in Ohiri, F.C., Verpoorte, R., and Baerheim Svendsen, A. 1983. The African *Strychnos* species and their alkaloids : A review. J. of Ethnopharmacology 9 : 167-223.

Tits, M., Desaive, C., Marnette, J.-M., Bassleer, R., and Angenot, L. 1984. Antimitotic activity of strychnopentamine, a bisindolic alkaloid. J. of Ethnopharmacology 12 : 287-292.

Toppel, G., Witte, L., and Hartmann, T. 1988. N-oxidation and degradation of pyrrolizidine alkaloids during germination of *Crotalaria scassellatii*. Phytochemistry 27 : 3757-3760.

Verpoorte, R. 1976. Pharmacognostical studies of some African Strychnos species. Ph.D. dissertation. University of London.

- _____. 1986. Method for the structure elucidation of alkaloids. Journal of Natural Products 49 : 1-25.
- Verpoorte, R., and Baerheim Svendsen, A. 1978. Alkaloids of *Strychnos dolichothyrsa* Gilg ex Onochie et Hepper. J.Pharm.Sci. 67 : 171-174.
- Verpoorte, R., Hylands, P.J., and Bisset, N.G. 1977. Carbon-13 n.m.r. spectroscopy of some *Strychnos* alkaloids. Organic Magnetic Resonance 9 : 567-571.
- Verpoorte, R., Kode, E.W., Doorne, H. van, and Baeheim Svendsen, A. 1978. Antimicrobial effect of the alkaloids from *Strychnos afzelii* Gilg. Planta Medica 33 : 237-242.
- Verpoorte, R., Beek, T.A. van, Thomassen, P.H.A.M., Aandewiel, J., and Svendsen, A.B. 1983. Screening of antimicrobial activity of some plants belonging to the Apocynaceae and Loganiaceae. J. of Ethnopharmacology 8 : 287-302.
- Verpoorte, R., Bohlin, L., Dwuma-Badu, D., Rolfsen, W., and Strombom, J. 1983. 11-Methoxy-macusine A. A new quaternary alkaloid from *Strychnos angolensis*. Journal of Natural Products 46 : 572-575.
- Waser, P.G. 1972. Chemistry and pharmacology of natural curare compounds. In J. Cheymol (section ed.), International encyclopedia of pharmacology and

- therapeutics (vol.1, section 14), pp.205-239.
Oxford : Pergamon Press. quoted in Ohiri, F.C.,
Verpoorte, R., and Baerheim Svendsen, A. 1983.
The African *Strychnos* species and their alkaloids
: A review. J. of Ethnopharmacology 9 : 167-223.
- Weeratunga, G., Goonetileke, A., Rolfsen, W., Bohlin, L.,
and Sandberg, F. 1984. Alkaloids in *Strychnos*
aculeata. Acta Pharm.Suec. 21 : 135-140.
- Wenkert, E., Andrew Cheung, H.T., and Gottlieb, H.E.
1978. Carbon-13 nuclear magnetic resonance spec-
troscopy of naturally occurring substances. 56.
Strychnos alkaloids. J.Org.Chem. 43 : 1099-1105.
- Zenk, M.H. 1980. Enzymatic synthesis of ajmalicine and
related indole alkaloids. Journal of Natural
Products 43 : 438-451.

APPENDIX




Figure 16 : Ultra-violet absorption spectrum of A-1

— = in MeOH
- - - = in MeOH+NaOH




Figure 17 : Infrared absorption spectrum of A-1 in KBr disc




Figure 18 : Mass spectrum of A-1




Figure 19 : ^1H Nuclear Magnetic Resonance spectrum
(300 MHz) of A-1 in CDCl_3




Figure 20 : ^{13}C Nuclear Magnetic Resonance spectrum
(22.5 MHz) of A-1 in CDCl_3




Figure 21 : Ultra-violet absorption spectrum of AA-1
in MeOH




Figure 22 : Infrared absorption spectrum of AA-1 in KBr disc




Figure 23 : Mass spectrum of AA-1






Figure 24 : ^1H Nuclear Magnetic Resonance spectrum
(80 MHz) of AA-1 in CDCl_3




Figure 25 : ¹H Nuclear Magnetic Resonance spectrum,
(300 MHz) of AA-1 in CDCl_3




Figure 26 : ^{13}C Nuclear Magnetic Resonance spectrum
(22.5 MHz) of AA-1 in CDCl₃

VITA

Mr. Uthai Sotanaphun was born on 3rd January, 1965 in Bangkok, Thailand. He obtained a B.Sc. in Pharm. from the Faculty of Pharmaceutical Science, Chulalongkorn University in 1988.